



ORIGINAL ARTICLE

**REHABILITATION OF ATROPHIC POSTERIOR MAXILLA WITH EXTRASINUS
IMPLANTS COMBINED WITH ANTERIOR STANDARD IMPLANTS**

Levon Khachatryan DDS, PhD ^{1*}, Grigor Khachatryan MD, PhD ², Edgar Karapetyan ³, Vardges Sahakyan ⁴

¹ Maxillofacial surgeon, Head of the Medical Center MIM (MC Maxillofacial and Plastic Surgery), Armenia

² Associate Professor of Department of Oral and Maxillofacial Surgery, Yerevan State Medical University after M. Heratsi, Armenia, Head of the Medical Center MIM (Medical Center of Maxillofacial and Plastic Surgery), Armenia

³ Maxillofacial surgeon, Medical Center MIM (Medical Center of Maxillofacial and Plastic Surgery), Armenia

⁴ Prosthodontist Medical Center MIM (Medical Center of Maxillofacial and Plastic Surgery), Armenia

**Corresponding author: Levon Khachatryan, Head of the Medical Center MIM (Medical Center of Maxillofacial and Plastic Surgery), Armenia;
e-mail: info@mim.am*

Received: Nov. 26, 2022; **Accepted:** Dec. 27, 2022; **Published:** Jan. 15, 2023

Abstract

Aims: To evaluate treatment success at dental implants placement in the tuber regions and zygomatic bone of the maxilla in patients who had edentulous atrophic maxilla.

Patients and Methods: The present retrospective study aimed at investigating the 5-year clinical treatments outcomes 28 patients with implants placement in the tuber regions and zygomatic bone of the maxilla.

Results: No intra-operative or immediate post-operative complications were noted. After 5-6 months implants healing time evaluation of CT scan revealed no radiolucency around the implants. Implants placed in the tuber and zygoma areas of the maxilla demonstrated to integrate normally, implants show survival rates (97%) after 5 years.

Conclusions: With proper case selection, correct indication and knowledge of the surgical technique, the dental implants placement in the tuber regions and zygomatic bone of the maxilla offers advantages in the rehabilitation of

Keywords: atrophic maxilla, maxillary sinus, tuber regions, zygomatic bone

Introduction

Dental implants are now commonly used for replacing missing teeth in various clinical situations. Infections, trauma during dental extraction, remodeling of alveolar bone after tooth extraction create localized defects on the bone, affecting its height and width, and consequently, influence the dental implant placement.

Bone resorption often making it impossible to place conventional dental implants in the posterior maxilla¹⁻⁴.

To date, there is no conclusive evidence in the literature on the superiority of one technique over the others in terms of prosthetic or implant success. The decision for either of the options, therefore, depends

upon patient factors, and ultimately, the expertise and skill of the clinician.

The treatment options for implant rehabilitation of atrophic maxilla can be broadly classified into two categories:

1. Augmentation of the bony defect.
2. Modified implant designs for specific conditions.

Many different surgical procedures have been developed to increase local bone volume in deficient anatomical regions, including total/segmental bone onlays and grafting of the maxillary sinus with autogenous bone and/or bone substitute⁵⁻⁸.

Any of these procedures requires considerable surgical expertise and has its own advantages, limits, surgical risks and complications involving biological and financial costs⁹⁻¹².

Sinus graft procedures whether autogenous or allogeneous, carries with it a risk of complications that include the harvesting procedure itself (for autogenous grafts) and the possibility of graft infection, poor flap closure, dehiscence and resorption of the graft^{13, 14}.

Many patients seeking treatment with osseointegrated implants meet the situation of severe resorption of alveolar bone and often they do not want pass thru a reconstructive surgery (increases morbidity, hospitalization, increases the treatment time, costs, surgical risks and other). Modern tendencies of dental implantation are aimed at minimizing surgical trauma and reducing the time for rehabilitation of patients. As well different alternative methods have been proposed, such as implants placed in specific anatomical areas like the pterygoid region, the tuber or the zygoma.

Dental implants placement in the tuber region and zygoma bone of the maxilla is one way to overcome the problem of insufficient bone volume for routine implant surgery in the posterior maxilla due to severe resorption of jawbone and an extensive enlargement of the maxillary sinus.

The aim of this study was to evaluate treatment success at dental implants placement in the tuber regions and zygomatic bone of the maxilla in patients who had edentulous atrophic maxilla.

Patients and Methods

The present study aimed at investigating the 5-year clinical treatments outcomesa 28 patients (15 males

and 13 females, the age was 46 to 68 years, from 2016 to 2022), who had edentulous atrophic maxilla. Before the implantation, the patient was examined and a comprehensive examination and treatment plan was drawn up. Clinical, laboratory, radiological methods were used in the examination of patients. Preoperative radiographs including cone beam were obtained for initial screening and evaluation. The treatment plan includes detailed analysis of space for restoration, bone quantity and density, radiographic techniques, selection of number, diameter, and length of the implants, and occlusion. Implants were inserted using CT images and measures from the planning software. Software made correct implants' positions in accordance with anatomical structures. The surgical procedure included full thickness flap protocol. A crestal incision was made on the edentulous ridge and the full thickness flap was elevated and bone was exposed. Implant bed was prepared (sequential increase in diameter). The implant was inserted by wrench to the level of the margin of the implant bed in the ridge. The healing screw was installed to the implant. The flap was replaced back in place and fixed in position with a 4-0 Vicryl suture. The sutures were then removed after 1 weeks.

A total of 64 implants had been surgically placed in the tuber regions, 56 long implants placement in the zygomatic bone of the maxilla and 108 implants in adjacent areas to support fixed dental bridges. After the surgery the patients were performed a control CT scan to make sure the position of the implant in the bone corresponds the planned positions. The implants were inserted at the correct position, inclination, and depth as planned in the 3D software. Implants planning and placement in accordance with the prosthetic treatment plan, may bring significant benefits to prosthetic rehabilitation procedures. Prosthodontic treatment was performed 5-6 months after implants healing time, 24 patients had received implant fixed prostheses and 4 patients had received implant-bridge and hybrid denture that provided ideal facial balance and occlusion. The prosthetic indication was made according to each patient clinical condition in order to achieve the highest function and esthetic.

Assessment of masticatory function was made both subjectively and objectively. Masticatory performance was objectively evaluated by chewing

of a piece of colorchangeable chewing gum (Xylitol, Lotte, Tokyo, Japan) for 60 strokes. This method is easy, simple, and quick, with no need for bulky equipment, and it has advantages in stimulating a natural and stable act of chewing while still allowing complete recovery of the test item. Colorchangeable chewing gum has been applied in various fields. This gum base contains red, yellow, and blue dyes, citric acid and xylitol. With the progression of chewing, the color of the chewing gum turns from yellowish-green to red²⁴. Positive value indicating redness, and negative value indicating greenness.

The outcomes of the study collected both directly at the time of surgery (intraoperative complications, stability of the implant), and at the annual checkups (implant survival). Postoperative clinical and radiographic controls were made regularly, the criteria for implant success were assessed.

Outcome measures were: prosthesis success; implant success; complications and marginal bone levels. Postsurgical change in marginal bone levels was assessed by digital x-ray were taken immediately (base line for comparison) and 1,3,5 years post operatively.

Table 2: The mean marginal bone loss of implants after 5 years

The mean crestal bone loss	Number of tuber regions implants (n=64)	Number of zygoma bone regions implants(n=56)	Number of conventional implants (n=108)
	1.6 mm± 0.25	1.4 mm± 0.29	1.2mm± 0.17

Of the 228 implants placed in these 28 patients, 4 failed to osseointegrate and 5 after 5 years of loading, implants show survival rates (97%) after 5 years. Implants placed in the tuber region and zygoma bone of the maxilla demonstrated to integrate normally, with success and survival rates comparable to those obtained in case of implants placed in native bone.

The results showed that implant treatment is effective to improve patients' masticatory efficiency. The gum initially had a greenish color and became more-and-more reddish with the duration and intensity of chewing, and there is a strong correlation between color change and masticatory performance and ability.

This case report presents a combination of surgical and prosthetic solutions applied to a case of

Results

No intra-operative or immediate post-operative complications were noted (table 1).

Table 1: Complications of surgery

Complications	Number of patients (n=28)
membrane perforation	0
errors in the implants position	0
implants inclination	0
pain in the operation area	9
Swelling	14

After 5-6 months implant placement evaluation of radiographies revealed no radiolucency around the implants. All of the patients presented with healthy soft tissue. After 5 years implant placement the mean marginal bone level at tuber regions implants was situated on average 1.6 mm (n=64) from the abutment-fixture junction, the zygoma bone implants showed an average bone level of 1.4 mm (n=56). The average (standard deviation) marginal bone loss on conventional implants was 1.2 mm (n=108) (table 2).

oral implant rehabilitation in patients with edentulism and severely atrophic maxillae.

Case report

A 46 years old patient, presented to our clinic with a diagnosis of generalized periodontitis of the lower jaw, with edentulous atrophic maxilla. A comprehensive clinical and radiographic evaluation revealed advanced alveolar bone resorption rendering the prognosis of all lower teeth unfavorable. After the preliminary clinical examination, a treatment plan was defined that included the removal of all the teeth of the mandible and the installation of 8 dental implants in lower jaw, the installation of 1 dental implants in the tuber regions of the upper jawbone on both sides, the installation of 2 dental implants in zygomatic bone from both sides and

installation 4 dental implants in the area of 13, 14, 23, 24. 4 months after implant placement prosthetic restoration was fabricated and adjusted (Figures 1-6). The patient has been followed up for 5 years the restoration has remained functional.



Figure 1: Preoperative CT scan



Figure 2: CT scan after 1 month surgery



Figure 3: Intraoral view of abutments upper jaw before prosthetic reconstruction;

Figure 4: Intraoral view of abutments lower jaw before prosthetic reconstruction;

Figure 5: Clinical appearance after prosthetic rehabilitation with non-removable metal-ceramic prosthetic restoration



Figure 6: CT scan after prosthetic rehabilitation

Discussions

Bone grafting and sinus lifts are invasive procedures. In addition, they add complexity and increase the number of surgical phases required for implant therapy. These techniques pose a series of inconveniences, such as the need for multiple surgical interventions, the use of extraoral bone donor sites (e.g., iliac crest or skull) - with the

morbidity involved in surgery of these zones - and the long duration during which patients remain without rehabilitation during the graft consolidation and healing interval²⁵. These factors complicate patient acceptance of the restorative treatment and limit the number of procedures carried out. Sinus lifts procedure is one of the most common preprosthetic surgical procedures performed in dentistry today. However, the development and improvement of alternative methods for restoring the integrity of the dentition with an extreme degree of atrophy of the alveolar crest of the upper jaw is very urgent. In clinical practice it is becoming increasingly common for patients to demand therapies that offer a good final result while at the same time reduce costs, healing time and the temporary inability to work. In order to overcome such limitations, different therapeutic alternatives have been proposed, such as, short implants, or implants placed in specific anatomical areas like the pterygoid region, the tuber or the zygoma²⁶⁻³⁰. Any of these procedures requires considerable surgical expertise and has its own advantages, limits, surgical risks and complications involving biological and financial costs. The present study shows good clinical outcome with standard implants placed in the tuber regions and zygomatic bone of the maxilla using a two-stage procedure. This offers a more simplified treatment approach, a decrease in biological impact and a more comfortable post-surgical period for the patient thanks to a quicker recovery time. Implants through the atrophied upper jaw in the tuber regions and zygomatic bone of the maxilla are a good alternative to maxillary sinus lift and to bone grafts in patients with posterior atrophic maxillae. These methods allow:

1. Avoid bone grafting, which is used in conditions of atrophy of the upper jaw when installing conventional dental implants
2. To shorten the terms of rehabilitation: to produce a fully functional and aesthetic prosthesis

Implantation methods that were used in this work can provide a patient who is not ready for risky, expensive and multi-stage surgical treatment, the possibility of avoiding more traumatic (such as bone transplantation in the patient) and less predictable types of surgical intervention.

Carrying out a comparative analysis of the different approaches to the treatment of adentia in patients

with severe maxillary atrophy in the area of the maxillary sinus, we came to the conclusion that a reasonable combination of different techniques can be achieved in order to achieve the optimal result. Rehabilitation using implants in the tuber regions and zygomatic bone of the maxilla is a predictable technique, it does not lack in possible complications, and therefore, it should be reserved only to professionals with vast surgical experience, as it requires a long learning curve.

Conclusions

With proper case selection, correct indication and knowledge of the surgical technique, the dental implants placement in the tuber regions and zygomatic bone of the maxilla offers advantages in the rehabilitation of severely resorbed maxillae, implants show survival rates (97%) after 5 years.

These methods of implantation should be considered as an alternative approach to solving problems arising during prosthetics of the atrophied upper jaw.

Conflict of interest and financial disclosure: The author declares that he has no conflict of interest and there was no external source of funding for the present study. None of the authors have any relevant financial relationship(s) with a commercial interest.

Funding: This research did not receive any specific grant from funding agencies in the public, commercial, or not-forprofit sectors.

Ethics approval and consent to participate: This protocol was approved by the Clinical Research Ethics Committee Yerevan State Medical University after M. Heratsi (protocol N3 17.11.16) and in accordance with those of the World Medical Association and the Helsinki Declaration.

REFERENCES

1. Alfaro F.N. Bone grafting in oral implantology techniques and clinical applications Quintessence. 2006; 233 p.
2. Houry F., Antoun H., Missika P. Bone augmentation in oral implantology. Quintessence. 2007;435p.
3. Sharan A., Madjar D. Maxillary sinus pneumatization following extractions: A radiographic study. Int J Oral Maxillofac Implants 2008;23(1):48-56.
4. Boyne P., Cole M., Stringer D., Shafquat J. A technique for osseous regeneration of deficient edentulous maxillary ridges. J. Oral Maxillofac. Surg 1985; 43:87-91.
5. Sailer H.A. A new method of inserting endosseous implants in totally atrophic maxillae. J CMS.1989;17(7):299-305.
6. Tonetti M.S., Hämmerle C.H. European Workshop on Periodontology Group C. Advances in bone augmentation to enable dental implant placement: Consensus Report of the Sixth European Workshop on Periodontology. J ClinPeriodontol 2008; 35: 168-72.
7. Rosen A., Gynther G. Implant treatment without bone grafting in edentulous severely resorbed maxillas: A long-term followup study. J Oral MaxillofacSurg 2007; 65: 1010-1016.
8. Maló P, NobreMde A, Lopes I. A new approach to rehabilitate the severely atrophic maxilla using extramaxillary anchored implants in immediate function: a pilot study. J Prosthet Dent. 2008; 100(5): 354-366.
9. Chanavaz M. Sinus graft procedures and implant dentistry: a review of 21 years of surgical experience (1979-2000). Implant Dentistry.2000; 3: 197-203.
10. Tatum H. Maxillary and sinus implant reconstruction Dent. Clin. North America. 1986; 30: 207-229.
11. Misch C.E. The maxillary sinus lifts and sinus graft surgery. In: Contemporary implant Dentistry. St Louis: Mosby 1999; 469- 495.
12. Peleg M., Mazor Z., Chaushu G., Garg A.K. Sinus floor augmentation with simultaneous implant placement in the severely atrophic maxilla. J Periodontol 1988;69:1397-1403.
13. Boyne P.J., James R.A. Grafting of the maxillary sinus floor with autogenous marrow and bone. J Oral Surg 1980; 38: 613-616.

15. Block M.S., Kent J.N. Sinus augmentation for dental implants, the use of autogenous bone. //J Oral Maxillofac Surg 1997; 55:1281-1286.
16. Balshi T.J., Wolfinger G.J., Balshi S.F. Analysis of 356 pterygomaxillary implants in edentulous arches for fixed prosthesis anchorage. Int J Oral Maxillofac Implants 1999; 14: 398-406.
17. Bidra A.S., Huynh-Ba G. Implants in the pterygoid region: a systematic review of the literature. Int J Oral Maxillofac Surg. 2011; 40(8): 773-781.
18. Candel E., Peñarrocha D., Peñarrocha M. Rehabilitation of the atrophic posterior maxilla with pterygoid implants: a review. J Oral Implantol. 2012; 38: 461-466.
19. Graves S.L. The pterygoid plate implant: a solution for restoring the posterior maxilla. // Int J Periodontics Restorative Dent 1994; 14: 512-523.
20. Ridell A., Grondahl K., Sennerby L. Placement of branemark implants in the maxillary tuber region: Anatomical considerations, surgical technique and longterm results. Clinical Oral Implants Research 2009;20(1):94-98.
21. Jaime G. Rodríguez-Chessa, Sergio Olate, Henrique Duque Netto, Jamil Shibli, Márcio de Moraes, and Renato Mazzonetto. Treatment of atrophic maxilla with zygomatic implants in 29 consecutives patients Int J ClinExp Med. 2014; 7(2):426-430.
22. Miglioranca R.M., Coppede, A., Dias Rezende R.C. and de Mayo T. Restoration of the edentulous maxilla using extrasinus zygomatic implants combined with anterior conventional implants: A retrospective study. The International Journal of Oral & Maxillofacial Implants, 2011; 26:665-672.
23. Candel-Marti E., Carrillo-Garcia, C., Penarrocha-Oltra, D. and Penarrocha-Diago, M. Rehabilitation of atrophic posterior maxilla with zygomatic implants: Review. Journal of Oral Implantology 2012; 38:653-657
24. Malevez C., Abarca M., Durdu F., Daelemans P. Clinical outcome of 103 consecutive zygomatic implants: A 6-48 months follow-up study. Clin Oral Implants Res. 2004;115:18- 22
25. Komagamine Y., Kanazawa M., Minakuchi S., Uchida T., Sasaki Y. Association between masticatory performance using a colour-changeable chewing gum and jaw movement. J Oral Rehabil 2011;38:555-63.
26. Iturriaga M.T., Ruiz C.C. Maxillary sinus reconstruction with calvarium bone grafts and endosseous implants. J Oral Maxillofac Surg. 2004; 62(3):344-347.
27. José-Luis Sierra-Sánchez et al. Predictability of short implants (<10 mm) as a treatment option for the rehabilitation of atrophic maxillae. A systematic review. Med Oral Patol Oral Cir Bucal. 2016; 21(3): 392-402.
28. Reiser G.M. Implant use in the tuberosity, pterygoid, and palatine region anatomical and surgical considerations. In: Nevins M, Mellonig JT (eds) Implant therapy clinical approaches and evidence of success, 2nd edn. Quintessence Books, Chicago, 1998.
29. Prithviraj R.V., Harleen K.B. From maxilla to zygoma: A review on zygomatic implants. Journal of Dental Implants 2014;4(1):44-47
30. Galán-Gil S., Penarrocha-Diago M., Balaguer-Martínez J., Martí-Bowen E. Rehabilitation of severely resorbed maxillae with zygomatic implants: an update. Med Oral Patol Oral Cir Bucal. 2007;12(3):216-220

ՎԵՐԻՆ ԾՆՈՏԻ ՀԵՏԻՆ ԱՊԱՃՈՎ ՀԱՏՎԱԾԻ ՎԵՐԱԿԱՆԳՆՈՒՄ ԷՔՍՏՐԱՍԻՆՈՒՍԱՅԻՆ ԻՄՊԼԱՆՏՆԵՐՈՎ ՀԱՄԱԿՑՎԱԾ ՖՐՈՆՏԱԼ ՀԱՏՎԱԾՈՒՄ ՏԵՂԱԴՐՎԱԾ ՍՏԱՆԴԱՐՏ ԻՄՊԼԱՆՏՆԵՐԻ ՀԵՏ

Լևոն Խաչատրյան ¹, Գրիգոր Խաչատրյան ², Էդգար Կարապետյան ³, Վարդգես Սահակյան ⁴

¹ Դիմաձևոտային վիրաբույժ, MIM բժշկական կենտրոնի ղեկավար, Հայաստան

² Երևանի Մ. Հերացու անվան պետական բժշկական համալսարանի վիրաբուժական ստոմատոլոգիայի և դիմաձևոտային վիրաբուժության ամբիոնի ղոցենտ, MIM բժշկական կենտրոնի դիմաձևոտային և պլաստիկ վիրաբուժության կլինիկայի ղեկավար, Հայաստան

Khachatryan L. et al. (2023). Rehabilitation of atrophic posterior maxilla with extrasinus implants combined with anterior standard implants. Bulletin of Stomatology and Maxillofacial Surgery. Vol. 19 N1, 40-46.

DOI: 10.58240/1829006X-2023.19.1-40

³ Դիմաձևնոտային վիրաբույժ, МИМ Դիմաձևնոտային և պլաստիկ վիրաբուժության բժշկական կենտրոն, Հայաստան.

⁴ Օրթոպեդ, МИМ Դիմաձևնոտային և պլաստիկ վիրաբուժության բժշկական կենտրոն, Հայաստան

Ամփոփում

Նպատակը: Գնահատել ադենտիայով հիվանդների մոտ վերին ծնոտի թմբի և այտոսկրի շրջաններում տեղադրված իմպլանտների բուժման արդյունավետությունը:

Հիվանդներ և մեթոդներ: Ներկայիս հետահայաց ուսումնասիրությունը նպատակաուղղված է հետազոտելու 5-ամյա կլինիկական բուժման արդյունքները 28 հիվանդների մոտ, որոնց իմպլանտները տեղադրվել են վերին ծնոտի թմբի և այտոսկրի շրջաններում:

Արդյունքներ: Ներվիրահատական կամ անմիջապես հետվիրահատական բարդություններ չեն նշվել: 5-6 ամիս հետո իմպլանտների գնահատումը CT սկանավորումամբ բացահայտեց, որ տեղադրված իմպլանտները նորմալ ինտեգրվել են, իմպլանտների գոյատևման մակարդակը (97%) 5 տարի հետո:

Եզրակացություններ: Պատշաճ դեպքի ընտրության, ճիշտ ցուցումների և վիրաբուժական տեխնիկայի իմացության դեպքում իմպլանտների տեղադրումը վերին ծնոտի թմբի և այտոսկրի շրջաններում առավելություններ է տալիս վերականգնողական գործընթացում և լավ այլընտրանք է սինուս լիֆտինգ վիրահատությանը:

РЕАБИЛИТАЦИЯ АТРОФИРОВАННОГО ЗАДНЕГО ОТДЕЛА ВЕРХНЕЙ ЧЕЛЮСТИ ЭКСТРАСИНУСНЫМИ ИМПЛАНТАТАМИ В СОЧЕТАНИИ С СТАНДАРТНЫМИ ПЕРЕДНИМИ ИМПЛАНТАТАМИ

Левон Хачатрян ¹, Григор Хачатрян ², Эдгар Карапетян ³, Вардгес Саакян ⁴

¹ Челюстно-лицевой хирург, руководитель Медицинского центра МИМ, Армения

² Челюстно-лицевой хирург, Доцент кафедры челюстно-лицевой хирургии, Ереванский государственный медицинский университет им. М. Гераци, Заведующий клиникой челюстно-лицевой и пластической хирургии Медицинского центра МИМ, Армения

³ Челюстно-лицевой хирург Медицинского центра МИМ, Армения

⁴ Ортопед Медицинского центра МИМ, Армения

Резюме

Цель: Оценить эффективность лечения при установке дентальных имплантатов в области бугра и скуловой кости верхней челюсти у пациентов с атрофической адентией верхней челюсти.

Пациенты и методы: Настоящее ретроспективное исследование было направлено на изучение результатов 5-летнего клинического лечения 28 пациентов с установкой имплантатов в области бугра и скуловой кости верхней челюсти.

Результаты: Интраоперационных или ближайших послеоперационных осложнений не отмечено. Через 5-6 месяцев времени заживления имплантатов оценка с помощью компьютерной томографии не выявила рентгенопрозрачности вокруг имплантатов. Имплантаты, установленные в области бугра и скуловой кости верхней челюсти продемонстрировали нормальную интеграцию, приживаемость имплантатов (97%) через 5 лет.

Выводы: При правильном выборе случая, правильных показаниях и знании хирургической техники, установка дентальных имплантатов в области бугра и скуловой кости верхней челюсти дает преимущества в реабилитации больных сильно резорбированной верхней челюсти.